

Mr John Coles Bury Hill Landscape Supplies Ltd The Estate Office Old Bury Hill Westcott Nr Dorking Surrey, RH4 3JU

> 11th June 2025 Our Ref: TOHA/25/1846/2/SS

> > Your Ref: see below

Dear Sirs

Topsoil Analysis Report: Bury Hill Horsham Yard - GP10 Plus Topsoil (S)

We have completed the analysis of the soil sample recently collected, referenced *GP10 Plus Topsoil (S)*, and have pleasure reporting our findings.

The purpose of the analysis was to determine the suitability of the sample for general landscape purposes (trees, shrubs, amenity grass). In addition, this sample has been assessed to determine its compliance with the requirements of the British Standard for Topsoil (BS3882:2015 – Specification for Topsoil – Table 1, Multipurpose Topsoil).

This report presents the results of analysis for the sample collected from the production site on 22/05/2025 and it should be considered 'indicative' of the topsoil source. The report and results should therefore not be used by third parties as a means of verification or validation testing, waste designation purposes or for any project-specific application, especially after the topsoil has left the Bury Hill Landscape Supplies Ltd site.

SAMPLE EXAMINATION

The sample was described as a very dark greyish brown (Munsell Colour 10YR 3/2), slightly moist, friable, slightly calcareous SANDY LOAM with weakly developed, very fine to fine granular structure. The sample had a very low stone content and contained moderate proportions of organic fines and occasional woody fragments. No unusual odours or deleterious materials were observed.



Plate 1: Sample GP10 Plus Topsoil (S)

ANALYTICAL SCHEDULE

The sample was submitted to a UKAS and MCERTS accredited laboratory for a range of physical and chemical tests to confirm the composition and fertility of the soil, and the concentration of selected potential contaminants. The following parameters were determined:

- detailed particle size analysis ('5 sands', silt, clay);
- stone content (2-20mm, 20-50mm, >50mm);
- saturated hydraulic conductivity;
- pH and electrical conductivity values;
- exchangeable sodium percentage;
- major plant nutrients (N, P, K, Mg);
- organic matter content;
- C:N ratio;
- heavy metals (Sb, As, B, Ba, Be, Cd, Cr, Cu, Pb, Hg, Ni, Se, V, Zn);
- total cyanide and total (mono) phenols;
- speciated PAHs (US EPA16 suite);
- aromatic and aliphatic TPH (C5-C35 banding);
- benzene, toluene, ethylbenzene, xylene (BTEX);
- asbestos screen.

The results are presented on the attached Certificate of Analysis and an interpretation of the results is given below.

TOHA/25/1846/2/SS/Jun Page 2

RESULTS OF ANALYSIS

Particle Size Analysis and Stone Content

The sample fell into the *sandy loam* texture class. Further detailed particle size analysis found the sample to have a predominance of *medium sand* (0.25-0.50mm), followed by *fine sand* (0.15-0.25mm) and *very fine sand*, (0.05-0.15mm), with a reasonable amount of *silt* and *clay*. Topsoils such as this are prone to self-compaction when initially placed in a landscape environment, which can lead to limited drainage and poor aeration, particularly if the soil has no developed structure. Placing the topsoil to a maximum depth of 300mm can help reduce the impact, which is in line with BS3882:2015, section A.3.

The stone content of the sample was very low and, as such, stones should not restrict the use of the soil for general landscape purposes.

Saturated Hydraulic Conductivity

The saturated hydraulic conductivity test is designed to test the sample's drainage potential under a moderate degree of compaction. The saturated hydraulic conductivity of the sample was moderate (14 mm/hr) and would be considered suitable for general landscape purposes.

pH and Electrical Conductivity Values

The sample was strongly alkaline in reaction (pH 8.4), with a low calcium carbonate (lime) content (2.6%). This pH value would be considered suitable for general landscape purposes provided species with a wide pH tolerance or those known to prefer alkaline soils are selected for planting, turfing and seeding.

The electrical conductivity (salinity) value (water extract) was moderate, which indicates that soluble salts should not be present at levels that would be harmful to plants.

The electrical conductivity value by CaSO₄ extract (*BS3882* requirement) fell below the maximum specified value (3300 μS/cm) given in *BS3882:2015 – Table 1*.

Organic Matter and Fertility Status

The sample was well supplied with organic matter and all major plant nutrients. The extractable potassium level (1554 mg/l) slightly exceeded the maximum permissible value (1500 mg/l) given in BS3882:2015 – Table 1.

The C:N ratio of the sample was acceptable for general landscape purposes.

Potential Contaminants

With reference to BS3882:2015 - Table 1: Notes 3 and 4, there is a requirement to confirm levels of potential contaminants in relation to the topsoil's proposed end use. This includes human health, environmental protection and metals considered toxic to plants. In the absence of site-specific assessment criteria, the concentrations that affect human health have been compared with the residential with homegrown produce land use in the Suitable For Use Levels (S4ULs) presented in The LQM/CIEH S4ULs for Human Health Risk Assessment (2015) and the DEFRA SP1010: Development of Category 4 Screening Levels (C4SLs) for Assessment of Land Affected by Contamination – Policy Companion Document (2014).

Of the potential contaminants determined, none exceeded their respective guideline values.

Phytotoxic Contaminants

Of the phytotoxic (toxic to plants) contaminants determined (copper, nickel, zinc), none was found at levels that exceeded the maximum permissible levels specified in BS3882:2015 – Table 1.

TOHA/25/1846/2/SS/Jun Page 3

CONCLUSION

The purpose of the analysis was to determine the suitability of the sample for general landscape purposes (trees, shrubs, amenity grass). In addition, this sample has been assessed to determine its compliance with the requirements of the British Standard for Topsoil (BS3882:2015 – Specification for Topsoil – Table 1, Multipurpose Topsoil).

From the soil examination and subsequent laboratory analysis, the sample was described as a strongly alkaline, non-saline, slightly calcareous sandy loam with an adequate structure and very low stone content. The sample contained moderate reserves of organic matter and all major plant nutrients. The extractable potassium level was slightly high. Of the potential contaminants determined, none exceeded their respective guideline values.

To conclude, based on our findings, the topsoil represented by this sample would be considered suitable for general landscape purposes (trees, shrubs and amenity grass), provided species with a wide pH tolerance or those known to prefer alkaline soils are selected and the physical condition of the soil is satisfactory.

To minimise the risk of self-compaction and anaerobism, we recommend that this soil is not placed thicker than a maximum depth of 300mm.

The sample was largely compliant with the requirements of the British Standard for Topsoil (BS3882:2015 – Specification for Topsoil – Table 1, Multipurpose Topsoil), with the exception of the slightly elevated extractable potassium level. In this instance, this non-compliance may be considered minor when reviewed in the context of all the results and the proposed end use of the soil. Extractable potassium is particularly soluble and the level should fall over time when leached by rainfall or additional irrigation.

Soil Handling Recommendations

It is important to maintain the physical condition of the soil and avoid structural damage during all phases of soil handling (e.g. stockpiling, respreading, cultivating, planting, seeding or turfing). As a consequence, soil handling operations should be carried out when soil is sufficiently dry to be non-plastic (friable) in consistency.

It is important to ensure that the soil is not unnecessarily compacted by trampling or trafficking by site machinery, and soil handling should be stopped during and after heavy rainfall and not continued until the soil is friable in consistency. If the soil is structurally damaged and compacted at any stage during the course of soiling or landscaping works, it should be cultivated appropriately to relieve the compaction and to restore the soil's structure prior to any planting, turfing or seeding.

Further details on soil handling are provided in Annex A of BS3882:2015.

We hope this report meets with your approval and provides the necessary information. Please do not hesitate to contact the undersigned if we can be of further assistance.

Yours faithfully

Harriet MacRae BSc MSc

H.MacKae

Soil Scientist

Ceri SpearsBSc MSc MISoilSci
Senior Associate

For & on behalf of Tim O'Hare Associates LLP

TOHA/25/1846/2/SS/Jun Page 4

Client:	Bury Hill Landscape Supplies Ltd
Project:	Bury Hill Horsham Yard
Job:	Topsoil Analysis - BS3882:2015
Date:	11/06/2025
Job Ref No:	TOHA/25/1846/2/SS



_			,	TIM OTTAKE ASSOCIA
Sample Reference			GP10 Plus	SOIL & LANDSCAPE CONSULTA
		A	Topsoil (S)	
Clay (<0.002mm)	%	Accreditation UKAS	10	
Silt (0.002-0.05mm)	%	UKAS	19	
Very Fine Sand (0.05-0.15mm)	%	UKAS	17	
Fine Sand (0.05-0.15hilli)	%	UKAS	18	
Medium Sand (0.25-0.50mm)	%	UKAS	24	
Coarse Sand (0.50-1.0mm)	%	UKAS	9	
Very Coarse Sand (1.0-2.0mm)	%	UKAS	3	
Total Sand (0.05-2.0mm)	%	UKAS	71	
Texture Class (UK Classification)		UKAS	SL	
Stones (2-20mm)	% DW	GLP	1	
Stones (20-50mm)	% DW	GLP	0	
Stones (>50mm)	% DW	GLP	0	
			1	
Saturated Hydraulic Conductivity (m)	mm/hr	A2LA	14	
pH Value (1:2.5 water extract)	units	UKAS	8.4	
Calcium Carbonate	%	UKAS	2.6	
Electrical Conductivity (1:2.5 water extract)	uS/cm	UKAS	1084	
Electrical Conductivity (1:2 CaSO ₄ extract)	uS/cm	UKAS	3280	
Exchangeable Sodium Percentage	%	UKAS	6.6	
			1	
Organic Matter (LOI)	%	UKAS	7.8	
Total Nitrogen (Dumas)	%	UKAS	0.36	
C : N Ratio	ratio	UKAS	13	. (/,
Extractable Phosphorus Extractable Potassium	mg/l	UKAS	70 1554	A'AKJ
Extractable Potassium Extractable Magnesium	mg/l	UKAS UKAS	1554 163	
Evitacianie iviadilesimii	mg/l	UNAS	103	
Visible Contaminants: Plastics >2.00mm	%	UKAS	0	
Visible Contaminants: Sharps >2.00mm	%	UKAS	0	
Total Antimony (Sb)	mg/kg	MCERTS	1.7	~ T J ¬
Total Arsenic (As)	mg/kg	MCERTS	9	
Total Barium (Ba)	mg/kg	MCERTS	38	
Total Beryllium (Be)	mg/kg	MCERTS	0.4	
Total Cadmium (Cd)	mg/kg	MCERTS	0	
Total Chromium (Cr)	mg/kg	MCERTS	17	
Hexavalent Chromium (Cr VI)	mg/kg	MCERTS	< 1.8	
Total Copper (Cu)	mg/kg	MCERTS	19	
Total Lead (Pb)	mg/kg	MCERTS	26	
Total Mercury (Hg)	mg/kg	MCERTS	< 0.3	
Total Nickel (Ni) Total Selenium (Se)	mg/kg	MCERTS MCERTS	< 1.0	
Total Vanadium (V)	mg/kg mg/kg	MCERTS	33	
Total Zinc (Zn)	mg/kg	MCERTS	50	
Water Soluble Boron (B)	mg/kg	MCERTS	1.9	
Total Cyanide (CN)	mg/kg	MCERTS	< 1.0	
Total (mono) Phenols	mg/kg	MCERTS	< 1.0	
Naphthalene	mg/kg	MCERTS	< 0.05	
Acenaphthylene	mg/kg	MCERTS	< 0.05	
Acenaphthene	mg/kg	MCERTS MCERTS	< 0.05 < 0.05	
Fluorene Phenanthrene	mg/kg mg/kg	MCERTS	0.25	
Anthracene	mg/kg	MCERTS	< 0.05	
Fluoranthene	mg/kg	MCERTS	0.48	
Pyrene	mg/kg	MCERTS	0.41	
Benzo(a)anthracene	mg/kg	MCERTS	0.15	
Chrysene	mg/kg	MCERTS	0.20	
Benzo(b)fluoranthene	mg/kg	MCERTS	0.18	
Benzo(k)fluoranthene	mg/kg	MCERTS	0.07	
Benzo(a)pyrene	mg/kg	MCERTS	0.12	
Indeno(1,2,3-cd)pyrene	mg/kg	MCERTS	0.09	
Dibenzo(a,h)anthracene	mg/kg	MCERTS	< 0.05	
Benzo(g,h,i)perylene	mg/kg	MCERTS	0.11	
Total PAHs (sum USEPA16)	mg/kg	MCERTS	2.06	
Aliphatic TPH (C5-C6)	mg/kg	MCERTS	< 0.010	
Aliphatic TPH (C6-C8)	mg/kg	MCERTS	< 0.010	
Aliphatic TPH (C8-C10)	mg/kg	MCERTS	< 0.010	
Aliphatic TPH (C10-C12)	mg/kg	MCERTS	< 1.0	
Aliphatic TPH (C12-C16)	mg/kg	MCERTS	< 2.0	
Aliphatic TPH (C16-C21)	mg/kg	MCERTS	< 8.0	
Aliphatic TPH (C21-C35)	mg/kg	MCERTS	17	
Aliphatic TPH (C5-C35)	mg/kg	MCERTS	17	
Aromatic TPH (C5-C7)	mg/kg	MCERTS	< 0.010	
Aromatic TPH (C7-C8)	mg/kg	MCERTS	< 0.010	
Aromatic TPH (C8-C10)	mg/kg	MCERTS	< 0.020	
Aromatic TPH (C10-C12)	mg/kg	MCERTS	< 1.0	
Aromatic TPH (C12-C16)	mg/kg	MCERTS	< 2.0	
Aromatic TPH (C16-C21)	mg/kg	MCERTS	< 10	
Aromatic TPH (C21-C35)	mg/kg	MCERTS	< 10	
Aromatic TPH (C5-C35)	mg/kg	MCERTS	< 10	
Benzene	mg/kg	MCERTS	< 0.005	
Toluene	mg/kg	MCERTS	< 0.005	
Ethylbenzene	mg/kg	MCERTS	< 0.005	
p & m-xylene	mg/kg	MCERTS	< 0.008	
o-xylene	mg/kg	MCERTS	< 0.005	
Asbestos	ND/D	ISO17025	Not-detected	

H.MacRae

Harriet MacRae BSc MSc Soil Scientist

SL = SANDY LOAM

Visual Examination

The sample was described as a very dark greyish brown (Munsell Colour 10YR 3/2), slightly moist, friable, slightly calcareous SANDY LOAM with weakly developed, very fine to fine granular structure. The sample had a very low stone content and contained moderate proportions of organic fines and occasional woody fragments. No unusual odours or deleterious materials were

Results of analysis should be read in conjunction with the report they were issued with.

The contents of this certificate shall not be reproduced without the express written permission of Tim O'Hare Associates LLP